

OUTBREAK STATUS REPORT

Middlesex-London Health Unit

Last Updated: December 1, 2021 7:33 PM

Active

Premise	Type of Outbreak	Date Declared	Affected Area(s)	Date Resolved	Pathogen 1	Pathogen 2	Pathogen 3
Amica & Company Retirement	Enteric	Nov 26, 2021	Floors 3-8		Unknown		

Declared Over

Premise	Type of Outbreak	Date Declared	Affected Area(s)	Date Resolved	Pathogen 1	Pathogen 2	Pathogen 3
Elmwood Place	Respiratory	Nov 20, 2021	East Floor	Dec 1, 2021	Rhinovirus		
Parkwood Institute Mental Health Care Building	Enteric	Nov 19, 2021	H3	Nov 27, 2021	Unknown		

This information is to be used as a guide and is intended to help you prevent infections and outbreaks in your facility. The decision regarding who can work at a facility rests with that facility, but should take into account the need for adequate staffing levels and the risk of infection from staff working at multiple facilities. For more information please visit <u>https://www.healthunit.com/outbreak-resources</u>

Enteric Outbreaks

- Staff members/volunteers from facilities in outbreak should not work at your facility until at least 48 hours have elapsed since they last worked at the affected facility. This period can be modified if the causative agent is known.
- Staff members/volunteers who are experiencing gastrointestinal illness should not work at any facility until they have been symptom-free for at least 48 hours. Good hand hygiene should be reinforced with returning staff members/volunteers.
- Staff should change to a clean uniform between facilities and before leaving the affected facility. (Source Control of Gastroenteritis Outbreaks in Long-Term Care Homes: A Guide for Long-Term Care Homes and Public Health Unit Staff, 2013). https://www.healthunit.com/outbreak-resources

Clostridium difficile Outbreaks

• Provided that all required infection control practices are in place, staff can work at both affected and unaffected facilities without restrictions. Residents/patients with suspected or confirmed *Clostridium difficile*-associated diarrhea (CDAD) can be transferred within the health care system; however, the receiving facility must be notified and be able to comply with the recommended infection control precautions. *Clostridium difficile* is suspected in a resident/patient who develops diarrhea and has been in a facility experiencing a *Clostridium difficile* outbreak. Initiate Isolation rapidly and stool samples should be collected. (Source: Annex C: Testing, Surveillance and Management of Clostridium difficile In All Health Care Settings, January 2013). https://www.healthunit.com/outbreak-resources

Respiratory Outbreaks (non-influenza)

• Staff members/volunteers from facilities in outbreak should not work at your facility until at least 72 hours have elapsed since they last worked at the affected facility. For outbreaks attributable to Respiratory Syncytial Virus (RSV), Parainfluenza virus, Human metapneumovirus, and Adenovirus consider lengthening this time period to 5 days.

Staff members/volunteers who are experiencing respiratory symptoms should not work at any facility until 5 days from onset of symptoms or until symptoms have resolved. For outbreaks attributable to RSV, lengthen this time period to 8 days. Good hand hygiene should be reinforced with returning staff members/volunteers. (Source: A Guide to the Control of Respiratory Infection Outbreak in Long-Term Care Homes, Ministry of Health and Long-Term Care, November, 2015). http://www.health.gov.on.ca/en/pro/programs/publichealth/flu/docs/resp infectn_ctrl_guide_ltc_2015_en.pdf

Influenza Outbreaks

- The Health Unit recommends that you ask non-immunized staff/volunteers who are from facilities in outbreak, and who are not on an appropriate antiviral drug, to refrain from working at your facility until it has been at least 72 hours since they last worked at the affected facility.
- Staff members/volunteers who are experiencing respiratory/ influenza symptoms should not work at any facility until 5 days from onset of symptoms or until symptoms have resolved, whichever time period is shorter. Good hand hygiene should be reinforced with returning staff members/volunteers. (Source: A Guide to the Control of Respiratory Infection Outbreak in Long-Term Care Homes, Ministry of Health and Long-Term Care, November, 2015). http://www.health.gov.on.ca/en/pro/programs/publichealth/flu/docs/resp_infect_ctrl_guide_ltc_2015_en.pdf